


# Unrealized Health Potential: A Snapshot of Iowa



## UNREALIZED HEALTH POTENTIAL AMONG CHILDREN

Based on two important indicators of health, infant mortality and children's general health status, children in Iowa are not as healthy as they could be. The levels of health for most Iowa children fall short of levels for children in the most-advantaged subgroups in the state and across the country. This snapshot describes these gaps as well as the social factors that are linked with these differences in health.

### INFANT MORTALITY

Iowa ranks 5th among states based on the size of the gap in infant mortality by mother's education, when comparing the current overall state rate of 5.5 deaths per 1,000 live births with the lower rate—4.2 deaths per 1,000 live births—seen among infants born to the state's most-educated mothers. Even if Iowa achieved this lower rate overall, infant mortality in the state would still exceed the *national benchmark* of 3.2 deaths per 1,000 live births—the lowest infant mortality rate seen in any state among babies born to mothers with 16 or more years of schooling. In Iowa, infant mortality rates in every maternal education and racial or ethnic group did not meet the national benchmark.

## CHILDREN'S GENERAL HEALTH STATUS

Iowa ranks 10th among states based on the size of the gap in children's general health status by family income, when comparing the current overall rate of 11.8 percent of children in less than optimal health with the lower rate—7.1 percent—seen among children in higher-income families. Even if Iowa achieved this lower rate overall, the state's rate would still exceed the *national benchmark* for children's general health status of 3.5 percent—the lowest rate of less than optimal health seen in any state among children in families that both were higher income and practiced healthy behaviors. In Iowa, the general health status of children in every income, education and racial or ethnic group did not meet the national benchmark.

## SOCIAL FACTORS AFFECTING CHILDREN'S HEALTH


Social factors such as income, education and racial or ethnic group can greatly affect a child's health. This snapshot describes these factors and how they are linked with infant mortality and children's general health status in the state.



IOWA:  
**Social Factors Affecting Children's Health**

Health during childhood is powerfully linked with social factors such as the income and education levels of a child's family and his or her racial or ethnic group. This snapshot of children ages 17 years or younger in Iowa shows that:

- One third of Iowa's children live in poor or near-poor households, two fifths live in middle-income households and one fourth live in higher-income households.
- Approximately one third of children in Iowa live in households where no one has education beyond high school, one third live with at least one person who has attended but not completed college and one third live with at least one college graduate.
- Over four fifths of Iowa's children are non-Hispanic white, 6 percent are Hispanic and 4 percent are non-Hispanic black.



Prepared for the RWJF Commission to Build a Healthier America by the Center on Social Disparities in Health at the University of California, San Francisco.  
 Source: 2006 American Community Survey (for data on income and racial or ethnic group); 2005–2007 Current Population Survey (for education data).  
<sup>†</sup> Guidelines set by the U.S. government for the amount of income providing a bare minimum of food, clothing, transportation, shelter and other necessities. In 2006, the U.S. FPL was \$16,079 for a family of three and \$20,614 for a family of four.  
<sup>‡</sup> "Other" includes children in any other racial or ethnic group or in more than one group.


## IOWA: Gaps in Infant Mortality

Infant mortality rates<sup>1</sup>—a key indicator of overall health—vary by mother’s education and racial or ethnic group in Iowa.

- Compared with babies born to the most-educated mothers, babies born to mothers with less education appear to be more likely to die before reaching their first birthdays. The infant mortality rate among babies born to mothers with less than 12 years of education is approximately 2.5 times that for babies born to mothers with 16 or more years of schooling.

- Babies born to non-Hispanic white mothers appear to have lower infant mortality rates than babies born to mothers in other racial or ethnic groups, although the difference in rates between Hispanics and non-Hispanic whites is not statistically significant.

Comparing Iowa’s experience against the national benchmark<sup>2</sup> for infant mortality reveals unrealized health potential among Iowa babies across maternal education and racial or ethnic groups. Infants in every group could do better.



Prepared for the RWJF Commission to Build a Healthier America by the Center on Social Disparities in Health at the University of California, San Francisco.

Source: 2000-2002 Period Linked Birth/Infant Death Data Set.

1 The number of deaths in the first year of life per 1,000 live births.

2 The national benchmark for infant mortality represents the level of mortality that should be attainable for all infants in every state. The benchmark used here—3.2 deaths per 1,000 live births, seen in New Jersey and Washington state—is the lowest statistically-reliable rate among babies born to the most-educated mothers in any state.

† Defined as any other or unknown racial or ethnic group, including any group representing fewer than 3 percent of all infants born in the state during 2000-2002.

## IOWA: Gaps in Children's General Health Status


Within Iowa, children's general health status<sup>1</sup> varies by family income and education and by racial or ethnic group. Children in the least-advantaged groups typically experience the worst health, but even children in middle-class families appear to be less healthy than those with greater advantages.

- Children in poor families are 3.5 times as likely and children in near-poor families are twice as likely to be in less than optimal health as children in higher-income families.

- Children in households without a high-school graduate are four times as likely to be in less than optimal health as children living with someone who has completed some college.

- Hispanic children are 4.5 times as likely to be in less than optimal health as non-Hispanic white children.

Comparing Iowa's experience against the national benchmark<sup>2</sup> reveals unrealized health potential among Iowa children in every income, education and racial or ethnic group.



Prepared for the RWJF Commission to Build a Healthier America by the Center on Social Disparities in Health at the University of California, San Francisco.  
Source: 2003 National Survey of Children's Health.

<sup>1</sup> Based on parental assessment and measured as poor, fair, good, very good or excellent. Health reported as less than very good was considered to be less than optimal.  
<sup>2</sup> The national benchmark for children's general health status represents the level of health that should be attainable for all children in every state. The benchmark used here—3.5 percent of children with health that was less than very good, seen in Colorado—is the lowest statistically-reliable rate observed in any state among children whose families were not only higher income but also practiced healthy behaviors (i.e., non-smokers and at least one person who exercised regularly).

† Defined as any other or more than one racial or ethnic group, including any group with fewer than 3 percent of children in the state in 2003.